



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

mixtures of aqueous vapour with any of the gases, according to which, each gaseous body is uniformly diffused throughout the whole space, its particles repelling those of its own kind, but exerting no pressure on the particles of any other kind. He considers the fact that a given portion of air has its volume expanded by the addition of aqueous vapour, as being of itself a sufficient refutation of that theory. The author then takes occasion to discuss the question, whether aqueous vapour exists in the atmosphere in the state of mechanical mixture or of chemical solution, and argues in favour of the latter view of the subject.

In the concluding section, the author enters at large into the investigation of the method of ascertaining heights by barometric observations, and gives various tables to be used for that purpose.

---

April 3, 1845.

The MARQUIS OF NORTHAMPTON, President, in the Chair.

Ἀπόρροια, No. 2. "On the Epipolic Dispersion of Light; being a Supplement to a paper entitled 'On a case of Superficial Colour presented by a Homogeneous Liquid internally colourless.'" By Sir John Frederick William Herschel, Bart., F.R.S. &c.

The author inquires whether the peculiar coloured dispersion of white light intronitted into a solution of sulphate of quinine, is the result of an analysis of the incident light into two distinct species, or merely of a simple subdivision analogous to that which takes place in partial reflexion, as exemplified in the colours of thin plates. He endeavours to ascertain the laws which regulate this singular mode of dispersion, which for brevity he terms *epipolic*, on account of the proximity of the seat of dispersion to the intronitting surface of the fluid. It might have been expected that by passing the same incident beam successively through many such dispersive surfaces, the whole of the blue rays would at length be separated from it, and an orange, or red residual beam be left: but the author establishes, by numerous experiments, the general fact, that an *epipolical beam of light*, meaning thereby a beam which has been once transmitted through a quiferous solution, and undergone its dispersing action, is incapable of farther undergoing epipolic dispersion.

There were only two liquids, out of all those examined by the author, namely oil of turpentine and pyroxylic spirit, which, when interposed in the incident beam, act like the solutions of quinine in preventing the formation of the blue film: and the only solid in which the author discovered a similar power of epipolic dispersion, is the green fluor of Alston Moor, and which by this action exhibits at its surface a fine deep blue colour.

---